

Dual mapping between the antisymmetric tensor matter field and the Kalb-Ramond field

L. Gonzaga Filho^a M. S. Cunha^a R. R. Landim^b

^a*Grupo de Física Teórica - GFT, Universidade Estadual do Ceará - Av. Paranjana, 1700, CEP 60740-000, Fortaleza, Ceará, Brazil*

^b*Departamento de Física, Universidade Federal do Ceará - Caixa Postal 6030, CEP 60455-760, Fortaleza, Ceará, Brazil*

Abstract

In this work we present a dual mapping between the Kalb-Ramond and antisymmetric tensor matter (ATM) field actions. Our procedure shows that the correlation functions associated with both the Noether current and the topological current are equivalent.

Key words: Kalb Ramond, matter field, dual mapping

PACS: 11.30.-j, 11.90.+t

Kalb-Ramond fields first appeared as a tensorial generalization of vector gauge fields. These allow the construction of topological invariants in D -dimensional manifolds [1,2,3] and have an important role in dualization [4,5,6,7]. Such antisymmetric tensor fields are the key to generate mass for the vector gauge fields through the topological mass mechanism [8,9,10]. More recently one finds some applications in the so-called physics of extra dimension. It is also noteworthy that the Kalb-Ramond fields appear in effective theories of specific low energy superstring models, and may describe axion physics or torsion of a Riemannian manifold [11].

Different types of (second rank) antisymmetric tensor fields have been introduced by Avdeev and Chizhov in [12]. There, on the other hand, the action was constructed with a matter field rather than a gauge field exhibiting several interesting features. As showed later by Lemes *et al* [13], in a BRST framework, some specific model which have antisymmetric tensor matter (ATM) fields is

¹ renan@fisica.ufc.br

renormalizable to arbitrary perturbative orders. In a subsequent paper they further realized that the ATM field is a real component of a complex tensor field that satisfies a complex self-dual condition [14]. As shown by the authors this condition makes the ATM field massless.

In a more recent paper [16] we proposed a mechanism to generate mass to the ATM field which preserves the $U(1)$ symmetry. By this mechanism, a topological term is introduced via a complex vector field. In [17] we also analyzed the possibility to give mass to a ATM field through the Higgs mechanism. There, a scalar field is coupled to the ATM field and requiring parity conservation it is described as a doublet where one of its components is a pseudo-scalar. We also showed that a topological term for the ATM field can be also generated by spontaneous symmetry breaking.

In spite of a large variety of papers thereafter published about these two types of tensors, no one has reported a possible connection between them. In the present paper we will show the existence of a dual mapping between the Kalb-Ramond and ATM fields. We apply the dual mapping method developed by Fosco *et al.* [18] by making use of the complex self-dual condition and the $U(1)$ symmetry.

Let us consider the action for the ATM field written in terms of the complex self-dual tensor [14]

$$S(\varphi, \varphi^\dagger) = \int d^4x \partial_\mu \varphi^{\mu\nu} \partial^\rho \varphi_{\rho\nu}^\dagger, \quad (1)$$

where $\varphi_{\mu\nu} = T_{\mu\nu} + i\tilde{T}_{\mu\nu}$ is the complex anti-symmetric tensor field satisfying the complex self-dual condition $\varphi_{\mu\nu} = i\tilde{\varphi}_{\mu\nu}$ [14]. In order to show the dual mapping procedure we first linearize the derivatives of the kinetic term with the introduction of a complex field $b_\mu = c_\mu + id_\mu$. We can now write the action (1) in the form

$$S(\varphi, b) = \int d^4x \left(b_\mu^\dagger \partial_\rho \varphi^{\rho\mu} + b^\mu \partial_\rho \varphi^{\rho\mu\dagger} - b^\mu b_\mu^\dagger \right). \quad (2)$$

The on-shell equivalence of (1) and (2) can be obtained by eliminating b_μ through the equation of motion

$$\frac{\delta S}{\delta b^{\dagger\mu}} = \partial^\rho \varphi_{\rho\mu} - b_\mu = 0. \quad (3)$$

The Noether current related to the global $U(1)$ invariance of Eq. (2) is obtained as usual by replacing the normal derivative of the tensor field $\varphi_{\mu\nu}$ by its covariant derivative $D_\mu \varphi_{\rho\sigma} = \partial_\mu \varphi_{\rho\sigma} + is_\mu \varphi_{\rho\sigma}$, namely

$$J^\mu = i \left(b_\nu^\dagger \varphi^{\mu\nu} - b_\nu \varphi^{\mu\nu\dagger} \right). \quad (4)$$

where s_μ is an external source.

The generating functional $\mathcal{Z}[s_\mu]$ for current-current correlation is

$$\mathcal{Z}[s_\mu] = \int [D\varphi Db] e^{-i(S(\varphi, b) + \int d^4x s_\mu J^\mu)}, \quad (5)$$

where the invariant functional measure can be written as

$$[D\varphi Db] = \delta(\varphi_{\mu\nu} - i\tilde{\varphi}_{\mu\nu}) D\varphi_{\mu\nu} D\varphi_{\mu\nu}^\dagger Db_\mu Db_\mu^\dagger. \quad (6)$$

From gauge invariance of (5), it follows that

$$\mathcal{Z}[s_\mu] = \int [D\varphi Db] D\eta_\mu \delta[f_{\mu\nu}(\eta)] e^{-i(S(\varphi, b) + \int d^4x (s_\mu + \eta_\mu) J^\mu)}. \quad (7)$$

Here we have defined $f_{\mu\nu} = \varepsilon_{\mu\nu\rho\sigma} \partial^\rho \eta^\sigma$, $\eta_\mu = \partial_\mu \alpha$, and α is a scalar field.

With the following representation of the Dirac's delta functional

$$\delta[f_{\mu\nu}(\eta)] = \int DB_{\mu\nu} e^{-i \int d^4x B_{\mu\nu} \varepsilon^{\mu\nu\rho\sigma} \partial_\rho \eta_\sigma}, \quad (8)$$

where $B_{\mu\nu}$ is an anti-symmetric tensor field, Eq. (7) can be written as

$$Z[s_\mu] = \int [D\varphi Db] DB_{\mu\nu} D\eta_\mu e^{-i(S(\varphi, b) + \int d^4x (s_\mu + \eta_\mu) J^\mu + \int d^4x B_{\mu\nu} \varepsilon^{\mu\nu\rho\sigma} \partial_\rho \eta_\sigma)}. \quad (9)$$

Redefining η_μ by $\eta_\mu = v_\mu - s_\mu$, the external source s_μ decouples from the matter field, and we get

$$Z[s_\mu] = \int [D\varphi Db] DB_{\mu\nu} Dv_\mu e^{-i(S(\varphi, b) + \int d^4x v_\mu J^\mu + \int d^4x B_{\mu\nu} \varepsilon^{\mu\nu\rho\sigma} \partial_\rho (v_\sigma - s_\sigma))}. \quad (10)$$

We can now define a dual action for $B_{\mu\nu}$ as follows

$$e^{-iS_{\text{dual}}[B_{\mu\nu}]} = \int [D\varphi Db] Dv_\mu e^{-i(S(\varphi, b) + \int d^4x v_\mu J^\mu + \int d^4x B_{\mu\nu} \varepsilon^{\mu\nu\rho\sigma} \partial_\rho v_\sigma)}, \quad (11)$$

such that

$$\mathcal{Z}[s_\mu] = \int DB_{\mu\nu} e^{-i(S_{\text{dual}}[B_{\mu\nu}] - \int d^4x s_\mu \varepsilon^{\mu\nu\rho\sigma} \partial_\nu B_{\rho\sigma})}. \quad (12)$$

As we can see from Eq. (11), $B_{\mu\nu}$ is in fact a true gauge field with a gauge invariant action

$$S_{\text{dual}}[B_{\mu\nu}] = S_{\text{dual}}[B_{\mu\nu} + \partial_\mu \omega_\nu]. \quad (13)$$

The $U(1)$ global symmetry for the fields and the complex self-dual condition are the key ingredients to implement our proposed duality. Actually, it is a

well-known fact that the existence of a global symmetry is crucial to obtain dualities.

Let us underline here that the same generating functional given by the equations (5) and (12) written in terms of $\varphi_{\mu\nu}$ or $B_{\mu\nu}$ gives the same correlation function for the $U(1)$ current and the topological current, *i.e.*,

$$\langle J_{\mu_1}(x_1) J_{\mu_2}(x_2) \dots J_{\mu_n}(x_n) \rangle_{\varphi_{\mu\nu}} = \langle j_{\mu_1}^T(x_1) j_{\mu_2}^T(x_2) \dots j_{\mu_n}^T(x_n) \rangle_{B_{\mu\nu}}, \quad (14)$$

where

$$j_{\mu}^T = \epsilon^{\mu\nu\rho\sigma} \partial_{\nu} B_{\rho\sigma}, \quad (15)$$

and

$$J_{\mu} = i \left(\partial_{\rho} \varphi \rho \nu^{\dagger} \varphi_{\mu\nu} - \partial_{\rho} \varphi^{\rho\mu} \varphi_{\mu\nu}^{\dagger} \right) = 2(\partial_{\rho} \tilde{T}^{\rho\nu} T_{\mu\nu} - \partial_{\rho} T^{\rho\nu} \tilde{T}_{\mu\nu}). \quad (16)$$

Note that both j_{μ}^T and J_{μ} are axial currents. As a consequence, parity is preserved in this duality. It is worth mentioning that the dual mapping developed in [18] breaks the parity symmetry for that scalar model in three dimension.

In conclusion, in this paper we have obtained a dual mapping between the Kalb-Ramond and the ATM field actions. This kind of dualization was implemented by the requirement of complex self-duality. As we saw, the same correlation function for the $U(1)$ current and the topological current was obtained. As a consequence parity is preserved and it is free of axial anomalies.

Acknowledgment

We wish to thank H.R Christiansen for a critical reading of the manuscript. The Conselho Nacional de Desenvolvimento Científico e tecnológico-CNPq is gratefully acknowledged for financial support.

Dedicatory

R. R. Landim - This paper is dedicated to the memory of my wife, Isabel Mara.

References

- [1] V. I. Ogievetsky and I. V. Polubarinov. The notoph and its possible interactions. *Yad.Fiz*, 4(1-2):216–223, 1966.
- [2] M. Kalb and P. Ramond. Classical direct interstring action. *Phys. Rev. D*, 9(8):2273–2284, 1974.
- [3] A. S. Schwarz. *Topological Conf. Abstracts*. Baku Intern., 1987.
- [4] A. Smailagic and E. Spallucci. Duality of massive gauge invariant theories in arbitrary space-time dimension. *Phys. Rev. D*, 61(6):067701, 2000.
- [5] A. Smailagic and E. Spallucci. Wess-zumino terms and duality. *Phys. Lett. B*, 489(3-4):427–434, 2000.
- [6] E. Harikumar and M. Sivakumar. On the equivalence between topologically and non-topologically massive abelian gauge theories. *Mod. Phys. Lett. A*, 15(2):121–131, 2000.
- [7] R. Menezes, J. R. S. Nascimento, R. F. Ribeiro, and C. Wotzasek. On the dual equivalence of the self-dual and topologically massive p-form models. *Phys. Lett. B*, 564(3-4):261–268, 2003.
- [8] T. J. Allen, M. J. Bowick, and A. Lahiri. Topological mass generation in 3+1 dimensions. *Mod. Phys. Lett. A*, 6(7):559–571, 1991.
- [9] I. Oda and S. Yahikozawa. Topologically massive nonabelian gauge-theories in higher dimensions. *Phys. Lett. B*, 234(1-2):69–71, 1990.
- [10] R. R. Landim and C. A. S. Almeida. Topologically massive non-abelian bf models in arbitrary space-time dimensions. *Phys. Lett. B*, 504(1-2):147–151, 2001.
- [11] B. Mukhopadhyaya, S. Sen, and S. SenGupta Does a Randall-Sundrum Scenario Create the Illusion of a Torsion-Free Universe? *Phys. Rev. Lett.*, 89(12):121101-1–121101-4, 2002.
- [12] L. V. Avdeev and M. V. Chizhov. Antisymmetric tensor matter fields - an abelian model. *Phys. Lett. B*, 321(3):212–218, 1994.
- [13] V. Lemes, R. Renan, and S. P. Sorella. Algebraic renormalization of antisymmetric tensor matter fields. *Phys. Lett. B*, 344(1-4):158–163, 1995.
- [14] V. Lemes, R. Renan, and S. P. Sorella. $\Phi(4)(4)$ -theory for antisymmetric tensor matter fields in minkowski space-time. *Phys. Lett. B*, 352(1-2):37–42, 1995.
- [15] B. Geyer and D. Mulsch. The topological b-model with antisymmetric tensor matter fields. *Phys. Lett. B*, 535(1-4):349–357, 2002.
- [16] L. Gonzaga Filho, M. S. Cunha, and R. R. Landim. *Europhys. Lett.*, 69(2):184–188, 2005.

- [17] L. Gonzaga Filho, M. S. Cunha, C. A. S. Almeida, and R. R. Landim. Phys. Lett. B, 646(5-6):279–281, 2007.
- [18] C. D. Fosco, V. E. R. Lemes, L. E. Oxman, S. P. Sorella, and O. S. Ventura. Dual description of $U(1)$ charged fields in (2+1) dimensions. *Ann. Phys.*, 290(1):27–44, 2001.